



Validation of on-line slaughter checks as a pig welfare diagnostic tool

Grace A. Carroll¹, Laura A. Boyle², Alison Hanlon³, Kym Griffin⁴, Lisa Collins⁵, and Niamh E. O'Connell¹

¹ Institute for Global Food Security, Queens University Belfast, Northern Ireland Technology Centre, Malone Road, Belfast BT9 5HN, UK


² Animal & Grassland Research & Innovation Centre, Teagasc Moorepark, Fermoy, Co Cork, Republic of Ireland

³ School of Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Republic of Ireland

⁴ School of Biological Sciences, Queens University Belfast, 97 Lisburn Road, Belfast BT9 7BL, UK

⁵ School of Life Sciences, University of Lincoln, Brayford Pool, Lincoln, Lincolnshire, LN6 7TS, UK

Email: gcarroll05@qub.ac.uk



Queen's University
Belfast



Department of
**Agriculture,
Food and the Marine**
An Roinn
**Talmhaíochta,
Bia agus Mara**

Introduction

Traditionally, farm animal welfare has been assessed by examining system inputs (Velarde & Dalmau, 2012), including space requirements, enrichment and floor type. However, only the **potential** welfare state of the animal can be assessed using this method. In response to this issue, novel output-based welfare assessment methods are being explored. The output-based nature of meat inspection (MI) makes it a good candidate for use as an animal welfare assessment tool. The overall aim of this research was to assess the extent to which welfare measures recorded at meat inspection at the abattoir reflect welfare assessments made throughout the life of pigs.

Method


Data were collected between May 2013 and December 2014 from ten batches of pigs (N = 720) reared in conventional intensive housing. 50% of the tail length was docked within 24 hours of birth. Each animal was assessed at 7, 9 and 10 weeks of age (early life), and at 15 and 20 weeks of age (later life). At each timepoint pigs were assessed for;



Tail lesions (TL)



Skin lesions (SL)



Health issues (HI)

Tail lesions were scored using a 5 point scale (adapted from Kritas and Morrison, 2007) that ranged from 0 - "no evidence of tail biting" to 5 - "total loss of tail". Skin lesions were scored using a 5 point scale (adapted from Conte et al., 2012) that ranged from 0 - "no injuries" to 5 - "many very big, deep and red lesions covering the skin area".

Health issues (HI) which were recorded at each timepoint included lameness, bursitis, coughing, scouring, rectal prolapse, hernias and aural hematomas. Following slaughter at 21 weeks, each carcass was then scored for tail length (long \geq 6cm, short \leq 5cm), tail lesions, loin bruising and fresh (red) and old (non-red) skin lesions. Levels of carcass condemnation were also recorded. Using a one-way Kruskal-Wallis ANOVA, the carcass measures of pigs with tail lesions, skin lesions and health issues in 'early life' (present at least once at 7 and 9 weeks only), 'later life' (present at least once in 10, 15 and 20 weeks only) or 'whole life' (present at least once in both early and later life) were compared to their respective controls (C).

Discussion

Skin and tail

lesions acquired in both early and later life remain visible on the carcass. Healed skin lesions appear to reflect skin damage acquired on farm. The finding that fresh skin lesions did not differ between groups suggests that these lesions may have been acquired during the marketing process. No carcass welfare measures reflected health issues recorded on farm. These issues may be reflected in traditional MI findings. However, there was an insufficient number of carcass condemnations for this to be examined. In conclusion, these findings show that carcass based measures of skin and tail lesions can be used to detect welfare problems occurring during both the early (growing) and later (finishing) period of pigs lives.

Results

Pigs recorded as having tail lesions in early life ($P < 0.05$), later life ($P < 0.001$) or during whole life ($P < 0.001$) had more carcass tail lesions than respective controls (Figure 1). Pigs recorded as having tail lesions in later life ($P = 0.001$) and during whole life ($P < 0.001$), but not during early life, had shorter tails on the carcass than C pigs. Pigs recorded as having skin lesions in early life, later life ($P < 0.05$) and during whole life ($P < 0.001$) had significantly more healed (non-red) carcass skin lesions than C pigs (Figure 2). Fresh (red) skin lesions to the carcass did not differ between the welfare groups. Levels of loin bruising on the carcass did not differ between any of the experimental groups ($P > 0.05$). In addition, health issues recorded during any stage of life were not reflected in any carcass measures ($P > 0.05$).




Figure 1 Mean carcass tail lesion scores




Figure 2 Mean healed carcass skin lesion scores

Acknowledgements

Thanks to the Irish Department of Agriculture, Food and the Marine for funding this research

References

- Conte, S., Lawlor, P. G., O'Connell, N. & Boyle, L. A. 2012. Effect of split marketing on the welfare, performance, and carcass traits of finishing pigs. *Journal of animal science*, 90, 373-380.
- Hansen, L. L. Retrieved from http://qpc.adm.slu.se/5_Entire_Male_Pigs/page_15.htm.
- Kritas, S. K. & Morrison, R. B. 2007. Relationships between tail biting in pigs and disease lesions and condemnations at slaughter. *Veterinary Record*, 160, 149-152.
- Velarde, A. & Dalmau, A. 2012. Animal welfare assessment at slaughter in Europe: Moving from inputs to outputs. *Meat Science*, 92, 244-251.
- Welfare Quality®, 2009. Welfare Quality® Assessment Protocol for Pigs. Welfare Quality® Consortium, Lelystad, Netherlands, 119 pp.